

# Effectiveness of a Nonrinse, Alcohol-Free Antiseptic Hand Wash

Anoosh Moadab, BS\*  
Kathryne F. Rupley, BS†  
Peter Wadhams, DPM‡

This study evaluated the efficacy of a novel surfactant, allantoin, and benzalkonium chloride hand sanitizer using the US Food and Drug Administration's method for testing antiseptic hand washes that podiatric physicians and other health-care personnel use. The alcohol-free product, HandClens, was compared with an alcohol-based product, Purell. Independent researchers from the California College of Podiatric Medicine conducted the study using 40 volunteer students from the class of 2001. The results show that HandClens outperformed Purell and met the regulatory requirements for a hand sanitizer. Purell failed as an antimicrobial hand wash and was less effective than a control soap used in the study. (*J Am Podiatr Med Assoc* 91(6): 288-293, 2001)

In today's health-care environment, prudent hand-washing practices have been adopted to decrease the transmission of bacteria from person to person.<sup>1</sup> However, the conscientious health-care workers, podiatric physicians, and others who follow these guidelines have a greater risk than less conscientious workers of developing a contact dermatitis due to repetitive hand washing and glove changing.<sup>2-7</sup> It has been established that the damaged skin of nurses can carry a greater number of bacterial pathogens associated with nosocomial infections than can healthy, undamaged skin.<sup>2</sup>

The irony of the antiseptics practices' causing dermatologic changes was originally discovered by Halstead.<sup>8,9</sup> Halstead invented the surgical glove as a means of reducing the hand irritation associated with the antimicrobial agents being used at the time. Although hand-washing formulations have changed since Halstead's time, the contact dermatitis associated with antimicrobial agents has remained.<sup>10</sup>

\*Submitted during fourth year, California College of Podiatric Medicine, San Francisco. *Mailing address:* 375 Laguna Honda Blvd, Box 27, San Francisco, CA 94116.

†Submitted during fourth year, California College of Podiatric Medicine, San Francisco.

‡Diplomate, American Board of Podiatric Surgery; Research Coordinator, Woodward Laboratories, Inc, Los Alamitos, CA.

The need for a broad-spectrum, long-lasting antimicrobial hand-wash product that protects the natural skin barrier motivated the development of an alcohol-free formulation of surfactants, allantoin, and benzalkonium chloride (SAB) in a hand spritz.<sup>11</sup> In this study, the efficacy of this novel formulation was compared with that of an alcohol-containing hand sanitizer using the US Food and Drug Administration (FDA) protocol for health-care personnel antiseptic hand washes. Independent researchers from the California College of Podiatric Medicine conducted the study using 40 volunteer students from the class of 2001.

## Materials and Methods

### Test Solutions

For this study, two solutions were evaluated using the FDA protocol (21 CFR 333.470) for health-care personnel antiseptic hand washes.

The first solution was the SAB hand wash, HandClens (Woodward Laboratories, Inc, Los Alamitos, California). The active ingredient in this product is benzalkonium chloride (0.13% vol/vol). Other ingredients are water, hydroxypropylmethyl cellulose, propylene glycol, cocamidopropylbetaine, cocamido-

propylamine oxide, cetyl, trimethyl ammonium chloride, quaternium-12, imidazolidinyl urea, quaternium-15, allantoin, methyl paraben, propyl paraben, eucalyptol, methyl salicylate, and triethanolamine.

The second solution was alcohol-based Purell (GOJO Industries, Akron, Ohio). The active ingredient is ethyl alcohol (62% vol/vol). Other ingredients are isopropyl alcohol, water, emollients, and thickener.

## Subjects

Forty volunteer students from the class of 2001 at the California College of Podiatric Medicine participated in this study. All subjects completed a 7-day quarantine period of abstaining from antimicrobial products prior to testing. All subjects had fingernails no longer than 2 mm. To be eligible for inclusion, subjects could have no abrasions or open lesions on their wrists or hands, and the wearing of jewelry was not permitted. Each subject gave informed verbal and written consent before entering into the study.

## Protocol

Each subject was asked to perform a practice wash using the control soap, Dove (Lever Pond's, Toronto, Ontario, Canada). Then, 5 g of control soap and 15 mL of sterile phosphate buffer were used to wash the hands for 2 min. Subjects then rinsed their hands under tap water for 30 sec.

Each subject was given a baseline inoculation: 5 mL of *Serratia marcescens* was placed in the subject's cupped hands. The subject rubbed the hands for 45 sec and allowed the hands to air-dry for another 2 min. Next, subjects underwent gloving and sampling. Large powder-free polyethylene gloves, with 50 mL of sterile phosphate buffer in each glove, were placed on the hands, secured with a rubber band at the wrist, and massaged for 1 min (gloving). Finally, both gloves were drained for "glove juice," using a sterile technique, into a sterile sampling tube. Undiluted and diluted samples were placed on agar plates, consisting of a combination of lecithin and polysorbate 20, and stored at room temperature, or 25°C (sampling).

Each subject was given a control wash. Subjects were inoculated once again and given 5 g of control soap to wash with for 2 min. They rinsed for 30 sec and then underwent gloving and sampling.

Each subject was allowed ten hand washes with the appropriate test solution. The steps were similar to those in the inoculation procedure: 5 mL of test solution with 2 min of washing, then gloving, and sampling only on hand washes 1, 3, 7, and 10. For

hand washes 2, 4, 5, 6, 8, and 9, glove juice was drained down the sink and all other steps remained the same.

Each sample was placed on an agar plate immediately. An undiluted plate was made. For more accurate counting, a diluted plate of a 1:10,000 dilution for baseline, a 1:1,000 dilution for control, and a 1:100 dilution for the hand-wash trials were made. In order to inactivate the benzalkonium chloride, and to prevent further microbicidal activity, a combination of lecithin and polysorbate 20 was used as the agar media. For inactivation of the alcohol-based sanitizer, simple dilution in the phosphate buffer was sufficient.

A 2-min intermission was employed between each contamination/hand-wash cycle and the next contamination/hand-wash cycle. Subjects refrained from touching any items after the practice wash was initiated. After the final hand wash, subjects were given their choice of antimicrobial scrub or soap.

## Data Analysis

Statistical analysis was performed using a Student's *t*-test available with Microsoft Excel (Microsoft, Inc, Redmond, Washington). For the purpose of comparison, mean differences were considered statistically significant if the confidence value returned from the test was 0.05 (ie,  $P < .05$ ).

## Results

For this experiment, the efficacy of a novel SAB formulation, HandClens, *versus* an alcohol-based, established formulation, Purell, was evaluated in terms of immediate and residual disinfecting power. Data sheets were kept for all subjects. The number of colony-forming units seen on the lecithin and polysorbate 20 agar plates for both undiluted and diluted samples was recorded for baseline contamination, control soap, wash 1 (test solution), wash 3, wash 7, and wash 10. Data on the individual subjects are provided in Tables 1 and 2. The efficacy of a test solution was calculated as a reduction factor (RF) defined by the FDA as the difference between the  $\log_{10}$  of the colony-forming unit for baseline contamination (CFU<sub>B</sub>) and the  $\log_{10}$  of the colony-forming unit for washes (CFU<sub>W</sub>):

$$RF = \log_{10} \text{CFU}_B - \log_{10} \text{CFU}_W$$

The FDA requires a minimum reduction factor value of 2 after a single hand wash and a minimum reduction factor value of 3 after the tenth hand wash for a hand sanitizer to be considered an acceptable antiseptic formulation. Statistical analysis was performed using

**Table 1. Data Collected for Subjects in the HandClens Group**

Subject ID	Baseline	Control	Wash 1	Wash 3	Wash 7	Wash 10
1	450,000	830	275	23	7	160
2	590,000	1,200	120,000	8	3	1
3	770,000	51,000	80	6	1	1
4	#	1,600	10,400	312	26	4
5	610,000	800	80,000	1	2	1
6	90,000	400	170	1	2	54
7	1,100,000	2,000	228	800	5	25
8	50,000	11,000	500	102	52	294
9	170,000	800	112	18	1	17
10	10,000	59,000	800	1	27	7
11	3,200,000	800	640	1	1	1
12	120,000	800	280	2	4	1
13	480,000	800	8,000	3	1	2
14	1,840,000	800	260	1	1	1
15	960,000	1,520	1,400	115	27	54
16	1,260,000	1,460	74,800	2	2	1
17	1,030,000	5,000	103	2	3	2
18	440,000	23,000	15,200	1	2,200	1
19	690,000	5,000	49	1	3	10
20	2,960,000	68,000	7,300	68	12	37

Note: Numbers shown indicate the number of colony-forming units with the following dilutions: baseline, 1:10,000; control, 1:1,000; hand wash, 1:100.

#Unable to calculate due to baseline sample overgrowth (too numerous to count).

**Table 2. Data Collected for Subjects in the Purell Group**

Subject ID	Baseline	Control	Wash 1	Wash 3	Wash 7	Wash 10
21	760,000	688	4,000	7,200	23,900	23,500
22	50,000	400	800	12,600	8,200	160,000
23	140,000	800	5,900	23,000	7,000	13,200
24	220,000	1,600	400	800	3,900	400
25	400,000	4,000	800	2,100	2,700	9,000
26	460,000	400	1,600	5,100	10,000	4,900
27	400,000	4,500	9,000	12,000	40,000	28,000
28	130,000	2,400	270	7,600	5,500	26,400
29	400,000	2,200	15,100	3,100	6,800	13,200
30	1,080,000	4,700	464	6,000	9,200	5,800
31	180,000	5,200	3,500	7,200	8,000	12,000
32	1,000,000	6,100	5,200	3,400	40,000	8,700
33	1,980,000	3,700	6,100	6,600	19,100	27,200
34	1,100,000	10,800	17,200	13,100	16,100	9,200
35	1,800,000	11,200	2,400	6,100	9,500	28,000
36	50,000	300	9,400	6,800	14,400	6,600
37	820,000	15,200	28,800	12,600	11,600	35,200
38	670,000	96,000	2,400	4,800	10,600	6,400
39	400,000	9,600	17,200	88,000	80,000	#
40	1,120,000	172,000	27,200	23,200	2,800	19,600

Note: Numbers shown indicate the number of colony-forming units with the following dilutions: baseline, 1:10,000; control, 1:1,000; hand wash, 1:100.

#Unable to calculate due to baseline sample overgrowth (too numerous to count).

the Student's *t*-test, and the results are shown in Tables 3 and 4.

The results showed that both groups met the minimum requirement for the first hand wash, with an average reduction factor value of 2.6 for HandClens and 2.1 for Purell. Next, an overall trend of sustained disinfecting power was seen for HandClens, as demonstrated by reduction factor values of 2.6, 4.9, 5.0, and 4.9 for hand washes 1, 3, 7, and 10, respectively. These values not only met the first requirement, but surpassed the minimum expected persistent values. This is noticeable at the third hand wash, which exceeds the expected persistence by 1.9 log<sub>10</sub> units, and also at the seventh hand wash, as demonstrated by 2.0 log<sub>10</sub> units.

In contrast, Purell's performance diminished over time and hand washes, as illustrated by reduction factor values of 2.1, 1.8, 1.6, and 1.5 for hand washes 1, 3, 7, and 10, respectively. Clearly, these values began to plummet as early as the third hand wash (Fig. 1) and failed to meet FDA standards for an anti-

septic hand sanitizer. Indeed, by the tenth hand wash, Purell's disinfecting abilities did not meet the minimal requirements with a 1.5 log<sub>10</sub> value. In fact, the antimicrobial capacity of Purell by the tenth hand wash was 0.5 log<sub>10</sub> less than that of the control soap (Dove). Surprisingly, only a 0.1 log<sub>10</sub> difference was found between the disinfecting ability of the nonantimicrobial control soap and that of the alcohol-based antimicrobial Purell. The antimicrobial activity of the alcohol-based hand sanitizer was significantly less (wash 1, *P* < .001; washes 3, 7, and 10, *P* < .001) than that of the alcohol-free HandClens product.

## Discussion

The value of using an antimicrobial hand sanitizer with an acceptable disinfecting power defined by the FDA for podiatric physicians and other health-care workers has already been emphasized. A nonirritating sanitizer with residual activity is ideal to preserve the natural skin barrier. This study evaluated the

**Table 3. Log Reductions for Subjects Using HandClens**


	Control	Wash 1	Wash 3	Wash 7	Wash 10
Subject ID					
1	2.7	3.2	4.3	4.8	3.4
2	2.7	0.7	4.9	5.3	5.8
3	1.2	4.0	5.1	5.9	5.9
4	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>
5	2.9	0.9	5.8	5.5	5.8
6	2.4	2.7	5.0	4.7	3.2
7	2.7	3.7	3.1	5.3	4.6
8	0.7	2.0	2.7	3.0	2.2
9	2.3	3.2	4.0	5.2	4.0
10	-0.8	1.1	4.0	2.6	3.2
11	3.6	3.7	6.5	6.5	6.5
12	2.2	2.6	4.8	4.5	5.1
13	2.8	1.8	5.2	5.7	5.4
14	3.4	3.8	6.3	6.3	6.3
15	2.8	2.8	3.9	4.6	4.2
16	2.9	1.2	5.8	5.8	6.1
17	2.3	4.0	5.7	5.5	5.7
18	1.3	1.5	5.6	2.3	5.6
19	2.1	4.1	5.8	5.4	4.8
20	1.6	2.6	4.6	5.4	4.9
Statistics					
Average	2.2	2.6	4.9	5.0	4.9
SD	1.0	1.2	1.0	1.2	1.2
<i>t</i> -value	9.266	9.86	20.62	18.44	17.63
<i>df</i>	18.0	18.0	18.0	18.0	18.0
<i>P</i> value	<.001	<.001	<.001	<.001	<.001

<sup>a</sup>Unable to calculate due to baseline sample overgrowth (too numerous to count).

**Table 4. Log Reductions for Subjects Using Purell**

	Control	Wash 1	Wash 3	Wash 7	Wash 10
Subject ID					
21	3.0	2.3	2.0	1.5	1.5
22	2.1	1.8	0.6	0.8	-0.5
23	2.2	1.4	0.8	1.3	1.0
24	2.1	2.7	2.4	1.8	2.7
25	2.0	2.7	2.3	2.2	1.6
26	3.1	2.5	2.0	1.7	2.0
27	1.9	1.6	1.5	1.0	1.2
28	1.7	2.7	1.2	1.4	0.7
29	2.3	1.4	2.1	1.8	1.5
30	2.4	3.4	2.3	2.1	2.3
31	1.5	1.7	1.4	1.4	1.2
32	2.2	2.3	2.5	1.4	2.1
33	2.7	2.5	2.5	2.0	1.9
34	2.0	1.8	1.9	1.8	2.1
35	2.2	2.9	2.5	2.3	1.8
36	2.2	0.7	0.9	0.5	0.9
37	1.7	1.5	1.8	1.8	1.4
38	0.8	2.4	2.1	1.8	2.0
39	1.6	1.4	0.7	0.7	*
40	0.8	1.6	1.7	2.6	1.8
Statistics					
Average	2.0	2.1	1.8	1.6	1.5
SD	0.6	0.7	0.6	0.5	0.7
t-value	15.75	13.9	12.34	13.08	9.357
df	19.0	19.0	19.0	19.0	18.0
P value	<.001	<.001	<.001	<.001	<.001

\*Unable to calculate due to baseline sample overgrowth (too numerous to count).



**Figure 1.** The average log reduction for the HandClens group and the Purell group for control wash using Dove soap and washes 1, 3, 7, and 10.

novel SAB formulation HandClens and compared its efficacy with that of an alcohol-based leading brand, Purell. The novel SAB formulation not only effectively killed microbes after the first wash, but continued to do so at a maximal value. The extent of the reduction factor was limited by the baseline contamination.

The efficacy of HandClens may be attributed to its unique SAB formulation. The combination was hypothesized to complement the natural skin barrier and enhance its performance, whereas alcohol-based formulations cause a deterioration of skin over time and with repetitive use. Clearly, HandClens surpassed the minimum FDA standards for an antiseptic hand sanitizer. Most importantly, the proven efficacy of HandClens will promote its adoption as an adjunctive hand sanitizer for busy podiatric physicians and other health-care professionals. Users of this SAB formulation can be assured of its disinfecting residual power and complementary action on their epidermis. The next concern should be testing the develop-

ment of more SAB products in different vehicles, such as surgical scrubs and hand lotions.

**Acknowledgment.** Lolita Segura, Eric Fuller, DPM, Armen Hagopjanian, BS, and David Dardashti for their assistance on this project.

## References

1. BRYAN JL, COHRAN J, LARSON EL: Hand washing: a ritual revisited. *Crit Care Nurs Clin North Am* 7: 617, 1995.
2. LARSON EL, HUGHES CAN, PYREK JD, ET AL: Changes in bacterial flora associated with skin damage on hands of health care personnel. *Am J Infect Control* 26: 513, 1998.
3. WYSS M, ELSNER P, WUTHRICH B, ET AL: Allergic contact dermatitis from natural latex without contact urticaria. *Contact Dermatitis* 28: 154, 1993.
4. NATIONAL INSTITUTE FOR OCCUPATIONAL SAFETY AND HEALTH: *NIOSH Alert: Preventing Allergic Reaction to Natural Rubber Latex in the Workplace*, US Department of Health and Human Services, Atlanta, GA, 1977.
5. EDITORS: Study reveals high prevalence of chapped hands. *Hosp Infect Control* 16: 94, 1989.
6. MASO MJ, GOLDBERG DJ: Contact dermatoses from disposable glove use: a review. *J Am Acad Dermatol* 23: 733, 1990.
7. HEESE A, VAN HINTZENSTERN J, PETERS KP, ET AL: Allergic and irritant reaction to rubber gloves in medical health services: spectrum, diagnostic approach, and therapy. *J Am Acad Dermatol* 25: 831, 1991.
8. CARTWRIGHT FF: Lister, the man: Lister centenary conference. *Br J Surg* 54: 405, 1967.
9. HOBSON DW, WOLLER W, ANDERSON L, ET AL: Development and evaluation of a new alcohol-based surgical hand scrub formulation with persistent antimicrobial characteristics and brushless application. *Am J Infect Control* 26: 507, 1998.
10. LARSON E, FRIEDMAN C, COHRAN J, ET AL: Prevalence and correlates of skin damage on hands of nurses. *Heart Lung* 26: 1714, 1997.
11. DYER DL, GERENRAICH KB, WADHAMS PS: Testing a new alcohol-free hand sanitizer to combat infection. *AORN J* 68: 239, 1998.

Available at: <http://www.cdc.gov/niosh/latexalt.html>.